

## **Wikiprint Book**

**Title: DATASETS**

**Subject: TracMeteo - udg/ecomms/dataserver/catalog**

**Version: 116**

**Date: 07/01/2022 06:26:34 AM**

## Table of Contents

|  |          |
|--|----------|
| <b>DATASETS</b>                                    | <b>4</b> |
| WFDEI (provided by the EU-funded WATCH project)    | 4        |
| ?NCEP_ reanalysis1 (provided by NCEP/NCAR)         | 4        |
| ERA_interim (provided by ECMWF)                    | 4        |
| System4 (provided by ECMWF)                        | 4        |
| CFSv2 (provided by NCEP)                           | 4        |
| GloSea5 (provided by Met-Office)                   | 4        |
| SMHI-EC-EARTH_EUPORIAS hindcast (provided by SMHI) | 5        |

The **ECOMS UDG** collects and provides information (mainly at 6-hourly and/or daily resolution) for a reduced number of variables from a number of datasets (seasonal hindcasts, reanalysis and observations) obtained from different data providers. The following list of variables has been identified according to the user's needs, receiving feedback from EUPORIAS WP22 (climate information indices, CII), WP23 (impact models), WP21 (calibration and downscaling) and SPECS WP61 (pilot applications) and WP52 (calibration and downscaling). See the section on the [assessment of user's needs](#) for more details.

Note that the **R names** below correspond to the vocabulary names used in the [R data access package](#) for homogenization purposes. Note that, data homogenization and aggregation (i.e. daily means from 6h data) is only provided through the R data access package.

In order to specify the particular **temporal frequency/aggregation** available for the variables in the different datasets, the following codes are used in the table below: **6h** (6-hourly instantaneous data). **12h** (12-hourly instantaneous data). **24h** (24-hourly instantaneous data). **DM** (daily mean value). **DX** (daily maximum value). **DN** (daily minimum value). **DA** (daily accumulated data). **DAR** (accumulated since the initialization time  $t_{runtime}$ ) NOTE: The R package performs deaccumulation on a daily basis to match the standard definition. **fx** (static field)

In the table below, boldface codes (e.g. **6h**) indicate variables already available through the ECOMS UDG. **Italics>** are used for work in progress (variables to be included in the next update). **e** indicates that a variable exists in the original dataset but it is not planned to be included yet in ECOMS-UDG; **blanks** indicate that the variables do not exist in the original dataset. Codes ended by (\*) indicate variables which do NOT exist in the dataset, but are derived/approximated from other available ones through the [R data access package](#). For more details on the particular approximations used see the [conversion formulae](#). Variables ended by (#) indicate daily aggregated values obtained from the corresponding original 3-hourly data.

| R name                                   | Variable description                       | Reanalysis:  |                      |                      | Seasonal forecasting models: |                  |                  |                |                      |                  | Observations | Global_reanalysis_12 | Global_reanalysis_24 |
|--|--|--------------|----------------------|----------------------|------------------------------|------------------|------------------|----------------|----------------------|------------------|--------------|----------------------|----------------------|
|  |  | WFDEI        | NCER_reanalysis      | ERA_interim          | System_reanal_12             | System_reanal_24 | System_reanal_15 | CFSr2_seasonal | System_reanal_12     | System_reanal_24 |              |                      |                      |
| <i>Surface variables</i>                 |  |              |                      |                      |                              |                  |                  |                |                      |                  |              |                      |                      |
| tss                                      | Near-Surface Air temperature               | DM           | 6h                   | DM                   | 6h/DM                        | DM/DM            |                  | 6h             |                      |                  | DM           | DM                   |                      |
| tssmax                                   | Daily Maximum Near-Surface Air Temperature | <b>DX(#)</b> | 6h                   | DX                   | DX                           | DX               | DX               | DX             | 6h                   |                  | DX           | DX                   |                      |
| tssmin                                   | Daily Minimum Near-Surface Air Temperature | <b>DN(#)</b> | 6h                   | DN                   | DN                           | DN               | DN               | DN             | 6h                   |                  | DN           | DN                   |                      |
| tp                                       | Total precipitation amount                 | DA           | 6hA                  | DA                   | DAR                          | DA/MA            | DAR              | 6h             |                      |                  | DA           | DA                   |                      |
| psl                                      | Sea Level Pressure                         |              | 6h                   | DM                   | 6h                           | 6h               | 12h/DM           | 6h             |                      |                  | DM           | DM                   |                      |
| ps                                       | Surface air pressure                       | DM           |                      | e                    | 6h(*)                        |                  |                  | 6h             |                      |                  |              |                      |                      |
| vs                                       | Wind speed (at 10m)                        | DM           |                      | e                    | 6h(*)                        | e                | e                | 6h(*)          |                      |                  |              |                      |                      |
| tsps                                     | 2m Dewpoint Temperature                    |              |                      | e                    | 6h                           | e                |                  |                |                      |                  |              |                      |                      |
| hurs                                     | Surface (2m) specific humidity             | DM           | 6h                   | e                    | 6h(*)                        |                  |                  | 6h             |                      |                  |              |                      |                      |
| hurs                                     | Surface (2m) relative humidity             |              | 6h                   | e                    | 6h(*)                        |                  |                  | 6h(*)          |                      |                  |              |                      |                      |
| rsds                                     | Surface Downwelling Shortwave Radiation    | DA           | 6hA                  | e                    | DAR                          | e                | e                | e              |                      |                  | DA           | DA                   |                      |
| rls                                      | Net Longwave Surface Radiation             | DA           | 6hA                  | e                    | DAR                          | e                | e                | e              |                      |                  |              |                      |                      |
| sst                                      | Sea surface temperature                    |              |                      | DM                   | e                            | e                | e                | e              |                      |                  |              |                      |                      |
| uas                                      | Eastward Near-Surface Wind                 |              | 6h                   | e                    | 6h                           | e                | e                | 6h             |                      |                  |              |                      |                      |
| vas                                      | Northward Near-Surface Wind                |              | 6h                   | e                    | 6h                           | e                | e                | 6h             |                      |                  |              |                      |                      |
| wsamax                                   | Wind speed (at 10m)                        |              |                      | e                    | e                            | e                | e                | e              |                      |                  |              |                      |                      |
| wgust                                    | Wind gust                                  |              |                      | e                    | e                            | e                | e                | e              |                      |                  |              |                      |                      |
| mrso                                     | Total Soil Moisture Content                |              |                      |                      |                              |                  |                  | e              |                      |                  |              |                      |                      |
| rmros                                    | Surface runoff flux                        |              |                      |                      |                              | e                | e                | e              |                      |                  |              |                      |                      |
| rmro                                     | Total Runoff                               |              |                      |                      |                              | e                | e                | e              |                      |                  |              |                      |                      |
| ssro                                     | Sub-surface runoff rate                    |              |                      |                      |                              | e                | e                | e              |                      |                  |              |                      |                      |
| wcsl                                     | Water Content of Soil Layer                |              |                      |                      |                              | e                | e                | e              |                      |                  |              |                      |                      |
| psn                                      | Snowfall amount                            | DA           |                      |                      | e                            | e                | e                | e              |                      |                  |              |                      |                      |
| sd                                       | Snow Depth                                 |              |                      |                      | 24h                          | e                | e                | e              |                      |                  |              |                      |                      |
| <i>3D vars @ isobaric surface levels</i> |  |              |                      |                      |                              |                  |                  |                |                      |                  |              |                      |                      |
| ua                                       | Eastward Wind                              |              | 6h @ standard levels | DM @ standard levels | 12h @ standard levels        | e                | e                | e              | 6h @ standard levels |                  |              |                      |                      |
| va                                       | Northward Wind                             |              | 6h @ standard levels | DM @ standard levels | 12h @ standard levels        | e                | e                | e              | 6h @ standard levels |                  |              |                      |                      |
| zg                                       | Geopotential height                        |              | 6h @ standard levels | DM @ standard levels | 12h @ standard levels        | e                | e                | e              | 6h @ standard levels |                  |              |                      |                      |
| ta                                       | Air temperature                            |              | 6h @ standard levels | DM @ standard levels | 12h @ standard levels        | e                | e                | e              | 6h @ standard levels |                  |              |                      |                      |
| hus                                      | Specific humidity                          |              | 6h @ standard levels | DM @ standard levels | 12h @ standard levels        | e                | e                | e              | 6h @ standard levels |                  |              |                      |                      |
| <i>Static fields</i>                     |  |              |                      |                      |                              |                  |                  |                |                      |                  |              |                      |                      |
| zgs                                      | Surface geopotential height                |              |                      |                      | fx                           | e                | e                | e              |                      |                  |              |                      |                      |
| orog                                     | Topography                                 | e            | e                    | e                    | e                            | e                | e                | e              | fx                   | e                |              |                      |                      |
| lm                                       | Binary land mask                           | e            | e                    | e                    | e                            | e                | e                | e              | fx                   | e                |              |                      |                      |

@ standard Levels: 1000,850,700,500,300,200 mb, except for hus, which is not available at 200mb in some models

**Data Homogenization:** The different nature of the datasets, and the idiosyncratic naming and storage conventions often applied by the modelling centres, makes necessary an homogenization across datasets in order to implement a truly user-friendly toolbox for data access. To this aim, the [R package for data access](#) has been developed. Data homogenization is achieved through the creation of a common vocabulary. The particular variables of each dataset are then translated -and transformed if necessary- into the common vocabulary by means of a *dictionary*. Both features -vocabulary and dictionary- are described [here](#). In particular, some typical transformations performed by the `loadECOMS` interface are deaccumulation of initialization-accumulated variables to daily accumulated (i.e.: **DAr** --> **DA**) and scaling and/or offset of variables to match standard units (e.g. -273.15 for conversion K --> °C).

## DATASETS

### WFDEI (provided by the EU-funded WATCH project)

The WATCH-Forcing-Data-ERA-Interim: [?WFDEI](#) was produced post-WATCH using WFD methodology applied to ERA-Interim data. It is a meteorological forcing dataset extending into early 21st C (1979 ? 2012). Eight meteorological variables at 3-hourly time steps, and as daily averages, for the global land surface at 0.5° x 0.5° resolution.

### ?NCEP\_ reanalysis1 (provided by NCEP/NCAR)

A subset of predictors commonly used in statistical downscaling.

### ERA\_interim (provided by ECMWF)

A subset of predictors commonly used in statistical downscaling (daily means at 2° resolution). This information has been downloaded from the ECMWF's MARS, degraded to a common 2 degrees grid and post-processed computing daily means based on the original 6 hourly fields when required. Therefore, this dataset is a degraded and subset of the original ERA-Interim reanalysis dataset, which is freely available via [?ECMWF servers](#) at original resolution.

This dataset is used in the framework of different international initiatives (such as CORDEX-ESD and VALUE) in order to have a standard predictor dataset and to facilitate the work of the contributing downscaling groups. More information [?here](#).

### System4 (provided by ECMWF)

The [?System 4](#) seasonal forecasting system became operational in November 2011. The corresponding hindcast is archived in the Meteorological Archival and Retrieval System ([?MARS](#)), the main data repository at the ECMWF, as a collection of GRIB-1 files at 0.75° spatial resolution. The downloaded data has been exposed as three different virtual datasets:

- [?System4\\_seasonal\\_15](#): There are twelve initializations (hereafter called *runtimes*) per year (the first of January, February, ...), each with 15 members running for 7 months (hereafter called simply *times*). Period: 1981-2010.
- [?System4\\_seasonal\\_51](#): There are only four *runtimes* per year (the first of February, May, August and November), each with 51 members running for 7 months. Period: 1981-2010.
- [?System4\\_annual\\_15](#): There are four *runtimes* per year each with 15 members, but the forecasts run for 13 months. Period: 1981-2010.

A preliminary [validation report](#) produced in SPECS (milestone MS22) is available for precipitation (`System4_seasonal_15`). The reports for all datasets and variables will be produced after feedback with end-users.

### CFSv2 (provided by NCEP)

The [?CFS version 2](#) seasonal forecasting model became operational at NCEP in March 2011. The corresponding [?retrospective CFSv2](#) forecast dataset is stored in the [?NOMADS server](#) as a collection of GRIB-2 files at 1° spatial resolution. The downloaded data is exposed as a single virtual dataset:

- [?CFSv2\\_seasonal](#). There are four initializations (4 cycles) from every 5th day (thus providing on average 24 members per month) running for 9 months (see [?the CFSv2 member configuration](#) for more detailed information of member construction). Period: 1982-2010. **Note:** For better comparability with other hindcasts, the [R data access package](#) defines by default an ensemble of 15 members for each lead month and forecast season.

### GloSea5 (provided by Met-Office)

GloSea5 data have been exposed as two different virtual datasets through ECOMS-UDG, based on their different forecast extents, initializations and member configuration:

#### [?Glosea5\\_seasonal\\_12](#)

- 4 initializations per year: Feb, May, Aug and Nov
- 4 × 3 members per initialization
- Total of 168 runtimes and 130 forecast times per runtime (~4.5 months)
- Hindcast period: 1996-2010

#### [?Glosea5\\_seasonal\\_24](#)

- 2 initializations per year: May and Nov
- 8 × 3 members per initialization
- Total of 126 runtimes and 120 forecast times per runtime (~4 months)
- Hindcast period: 1992-2013

Members are defined from all combinations of the members and lagged runtimes. In particular, in UDG members are ordered considering all combinations of member × runtime, being member the factor that varies fastest. For instance, a call to `loadECOMS` using `members = 1:5` in `dataset = Glosea5_seasonal_12` will return, in this order, the following 5 members:

-Member1: [member1,runtime1]  
 -Member2: [member2,runtime1]  
 -Member3: [member3,runtime1]  
 -Member4: [member4,runtime1]  
 -Member5: [member1,runtime2]

#### **SMHI-EC-EARTH\_EUPORIAS hindcast (provided by SMHI)**

The downloaded data is exposed as a single virtual dataset:

- [?SMHI-EC-EARTH\\_EUPORIAS](#) hindcast. There is one initialization, corresponding to May and 15 members running for 4 months. Period: 1991-2012. **Note:** For better comparability with other hindcasts, the [R data access package](#) defines by default an ensemble of 15 members for each lead month and forecast season.

---

Visit the [UDG wiki](#) for additional groups and datasets provided by the UDG.

---