

The **ECOMS UDG** collects and provides information (mainly at 6-hourly and/or daily resolution, but also some monthly data) for a reduced number of variables from a number of datasets (seasonal hindcasts, reanalysis and observations) obtained from different data providers. The following list of variables has been identified according to the user's needs, receiving feedback from EUPORIAS WP22 (climate information indices, CIIs), WP23 (impact models), WP21 (calibration and downscaling) and SPECS WP61 (pilot applications) and WP52 (calibration and downscaling). See the section on the [assessment of user's needs](#) for more details.

Note that the **R names** below correspond to the vocabulary names used in the [R data access package](#) for harmonization purposes. Data harmonization and temporal aggregation (i.e. daily means from 6h data) is **only** provided through the R data access package.

The original **temporal frequency/aggregation** of the variables stored at the UDG, the following codes are used in the table below: **6h** (6-hourly instantaneous data). **12h** (12-hourly instantaneous data). **24h** (24-hourly instantaneous data). **DM** (daily mean value). **DX** (daily maximum value). **DN** (daily minimum value). **DA** (daily accumulated data). **DAr** (accumulated since the initialization time $t_{runtime}$) NOTE: The R package performs deaccumulation on a daily basis to match the standard definition. **fx** (static field). Similarly, the prefix M indicates monthly data in the following way: **MM** (monthly mean), **MA** (monthly accumulated).

In the table below, boldface codes (e.g. **6h**) indicate variables already available through the ECOMS UDG. **Italics** are used for work in progress (variables to be included in the next update). **e** indicates that a variable exists in the original dataset but it is not planned to be included yet in ECOMS-UDG; **blanks** indicate that the variable does not exist in the original dataset. Codes ended by **(*)** indicate variables which do NOT exist in the dataset, but are derived/approximated from other available ones through the [R data access package](#). For more details on the particular approximations used see the [?conversion formulae](#). Variables ended by **(#)** indicate daily aggregated values obtained from the corresponding original 3-hourly data.

R name	Variable description	Observations		Reanalysis:		Seasonal forecasting models:					
		WRFDE	WRFDE	ERA-Interim	System, reanalysis_1	System, reanalysis_2	System, reanalysis_3	CFSv2, reanalysis	System, reanalysis_1	System, reanalysis_2	
Surface variables											
tss	Near-Surface air temperature	DM	6h	DM	6h/DM	DM/DM		6h		DM	DM
tssmax	Daily Maximum Near-Surface Air Temperature	DX(6h)	6h	DX	DX	DX	DX	6h		DX	DX
tssmin	Daily Minimum Near-Surface Air Temperature	DN(6h)	6h	DN	DN	DN	DN	6h		DN	DN
tp	Total precipitation amount	DA	6hA	DA	DAr	DA/MA	DAr	6h		DA	DA
psl	Sea Level Pressure		6h	DM	6h	6h/MM	12h	6h		DM	DM
ps	Surface air pressure	DM		e	6h(*)			6h			
uss	Wind speed (at 10m)	DM		e	6h(*)	e	e	6h(*)			
ltps	2m Deepest Temperature			e	6h	e					
hus	Surface (2m) specific humidity	DM	6h	e	6h(*)			6h			
hurs	Surface (2m) relative humidity	e	6h	e	6h(*)			6h(*)			
rsds	Surface Downwelling Shortwave Radiation	DA	6hA	e	DAr	e	e			DA	DA
rls	Net Longwave Surface Radiation	DA	6hA	e	DAr	e	e				
sst	Sea surface temperature			DM	e	e	e				
uss	Eastest Near-Surface Wind		6h	e	6h	e	e	6h			
uss	Northward Near-Surface Wind		6h	e	6h	e	e	6h			
ussmax	Wind speed (at 10m)			e	e	e	e				
ugust	Wind gust			e	e	e					
msso	Total Soil Moisture Content							e			
mtos	Surface runoff flux					e	e	e			
mrro	Total Runoff					e	e				
ssso	Sub-surface runoff rate					e	e				
wcsl	Water Content of Soil Layer					e	e				
psn	Snowfall amount	DA			e	e					
sd	Snow Depth				24h	e	e				
30 years @ isobaric surface levels											
ua	Eastward Wind		6h @ standard levels	DM @ standard levels	12h @ standard levels	e	e	e		6h @ standard levels	
ua	Northward Wind		6h @ standard levels	DM @ standard levels	12h @ standard levels	e	e	e		6h @ standard levels	
zg	geopotential height		6h @ standard levels	DM @ standard levels	12h @ standard levels	e	e	e		6h @ standard levels	
ta	Air temperature		6h @ standard levels	DM @ standard levels	12h @ standard levels	e	e	e		6h @ standard levels	
hus	Specific humidity		6h @ standard levels	DM @ standard levels	12h @ standard levels	e	e	e		6h @ standard levels	
Static fields											
zgp	Surface geopotential height				fx	e	e	e			
orog	Topography	e	e	e	e	e	e	e	fx	e	
lm	Binary land mask	e	e	e	e	fx	e	e	fx	e	

@ standard Levels: 1000,850,700,500,300,200 mb, except for hus, which is not available at 200mb in some models

Data Harmonization: The different nature of the datasets, and the idiosyncratic naming and storage conventions often applied by the modelling centres, makes necessary an harmonization across datasets in order to implement a truly user-friendly toolbox for data access. To this aim, the [R package for data access](#) has been developed. Data harmonization is achieved through the creation of a common vocabulary. The particular variables of each dataset are then translated -and transformed if necessary- into the common vocabulary by means of a *dictionary*. Both features -vocabulary and dictionary- are described [here](#). In particular, some typical transformations performed by the `loadECOMS` interface are deaccumulation of initialization-accumulated variables to daily accumulated (i.e.: **DAr** --> **DA**) and scaling and/or offset of variables to match standard units (e.g. -273.15 for conversion K --> °C).

DATASETS

WFDEI (provided by the EU-funded WATCH project)

The WATCH-Forcing-Data-ERA-Interim: [?WFDEI](#) was produced post-WATCH using WFD methodology applied to ERA-Interim data. It is a meteorological forcing dataset extending into early 21st C (1979 ? 2012). Eight meteorological variables at 3-hourly time steps, and as daily averages, for the global land surface at 0.5° x 0.5° resolution.

?NCEP_ reanalysis1 (provided by NCEP/NCAR)

A subset of predictors commonly used in statistical downscaling.

ERA_interim (provided by ECMWF)

A subset of predictors commonly used in statistical downscaling (daily means at 2° resolution). This information has been downloaded from the ECMWF's MARS, degraded to a common 2 degrees grid and post-processed computing daily means based on the original 6 hourly fields when required. Therefore, this dataset is a degraded and subset of the original ERA-Interim reanalysis dataset, which is freely available via [?ECMWF servers](#) at original resolution.

This dataset is used in the framework of different international initiatives (such as CORDEX-ESD and VALUE) in order to have a standard predictor dataset and to facilitate the work of the contributing downscaling groups. More information [?here](#).

System4 (provided by ECMWF)

The [?System 4](#) seasonal forecasting system became operational in November 2011. The corresponding hindcast is archived in the Meteorological Archival and Retrieval System ([?MARS](#)), the main data repository at the ECMWF, as a collection of GRIB-1 files at 0.75° spatial resolution. The downloaded data has been exposed as three different virtual datasets:

- [?System4_seasonal_15](#): There are twelve initializations (hereafter called *runtimes*) per year (the first of January, February, ...), each with 15 members running for 7 months (hereafter called simply *times*). Period: 1981-2010.
- [?System4_seasonal_51](#): There are only four *runtimes* per year (the first of February, May, August and November), each with 51 members running for 7 months. Period: 1981-2010.
- [?System4_annual_15](#): There are four *runtimes* per year each with 15 members, but the forecasts run for 13 months. Period: 1981-2010.

A preliminary [validation report](#) produced in SPECS (milestone MS22) is available for precipitation (`System4_seasonal_15`). The reports for all datasets and variables will be produced after feedback with end-users.

CFSv2 (provided by NCEP)

The [?CFS version 2](#) seasonal forecasting model became operational at NCEP in March 2011. The corresponding [?retrospective CFSv2](#) forecast dataset is stored in the [?NOMADS server](#) as a collection of GRIB-2 files at 1° spatial resolution. The downloaded data is exposed as a single virtual dataset:

- [?CFSv2_seasonal](#). There are four initializations (4 cycles) from every 5th day (thus providing on average 24 members per month) running for 9 months (see [?the CFSv2 member configuration](#) for more detailed information of member construction). Period: 1982-2010. **Note:** For better comparability with other hindcasts, the [R data access package](#) defines by default an ensemble of 15 members for each lead month and forecast season.

GloSea5 (provided by Met-Office)

GloSea5 data have been exposed as two different virtual datasets through ECOMS-UDG, based on their different forecast extents, initializations and member configuration:

[?Glosea5_seasonal_12](#)

- 4 initializations per year: Feb, May, Aug and Nov
- 4 × 3 members per initialization
- Total of 168 runtimes and 130 forecast times per runtime (~4.5 months)
- Hindcast period: 1996-2010

[?Glosea5_seasonal_24](#)

- 2 initializations per year: May and Nov
- 8 × 3 members per initialization
- Total of 126 runtimes and 120 forecast times per runtime (~4 months)
- Hindcast period: 1992-2013

Members are defined from all combinations of the members and lagged runtimes. In particular, in UDG members are ordered considering all combinations of member × runtime, being member the factor that varies fastest. For instance, a call to `loadECOMS` using `members = 1:5` in `dataset = Glosea5_seasonal_12` will return, in this order, the following 5 members:

-Member1: [member1, runtime1]

-Member2: [member2, runtime1]

-Member3: [member3, runtime1]

-Member4: [member4, runtime1]

-Member5: [member1, runtime2]

SMHI-EC-EARTH_EUPORIAS hindcast (provided by SMHI)

The downloaded data is exposed as a single virtual dataset:

- [?SMHI-EC-EARTH_EUPORIAS](#) hindcast. There is one initialization, corresponding to May and 15 members running for 4 months. Period: 1991-2012. **Note:** For better comparability with other hindcasts, the [R data access package](#) defines by default an ensemble of 15 members for each lead month and forecast season.

Visit the [UDG wiki](#) for additional groups and datasets provided by the UDG.
